

Scarlet Fever and Strep A

What you need to know

What is Scarlet Fever?

Scarlet fever is usually a mild illness, but it is highly infectious.

While for the vast majority of children this is a mild illness, early treatment with antibiotics will help avoid more serious illness such as pneumonia or a bloodstream infection.

If your child has scarlet fever, keep them at home until at least 24 hours after the start of antibiotic treatment to avoid spreading the infection to others.

Scarlet fever is caused by bacteria called group A streptococci. These bacteria also cause other respiratory and skin infections such as strep throat and impetigo.

In very rare occasions, the bacteria can get into the bloodstream and cause an illness called invasive Group A strep (iGAS). While still uncommon, there has been an increase in invasive Group A strep cases this year, particularly in children under 10.

Scarlet Fever symptoms include:

- Sore throat
- Headache
- Fever
- A fine pinkish or red body rash with a sandpapery feel. On darker skin, the rash can be more difficult to see, but will still have a sandpaper feel



Contact NHS 111 or your GP if you suspect your child has scarlet fever.

Scarlet Fever:

Your questions answered

Keeping children with Scarlet Fever at home will reduce the spread of infection. Early treatment will help reduce symptoms and the risk of more serious illness.


Good hand and respiratory hygiene are important for stopping the spread of many bugs. Teach your child to:

- wash their hands properly with soap for 20 seconds
- use a tissue to catch coughs and sneezes
- keep away from others when feeling unwell



Is this a new strain?

Currently, there is no evidence that a new strain is circulating. The increase is most likely related to high amounts of circulating bacteria and social mixing.



What should I look out for?

Look out for the following symptoms, and call 111 if you are concerned:

- your child is getting worse
- your child is feeding or eating much less than normal
- your child has had a dry nappy for 12 hours or more or shows other signs of dehydration
- your baby is under 3 months and has a temperature of 38°C, or is older than 3 months and has a temperature of 39°C or higher
- your baby feels hotter than usual when you touch their back or chest, or feels sweaty
- your child is very tired or irritable



When should I call 999 or go to A&E?

Call 999 or go to A&E if any of the following:

- your child is having difficulty breathing – you may notice grunting noises or their tummy sucking under their ribs
- there are pauses when your child breathes
- your child's skin, tongue or lips are blue
- your child is floppy and will not wake up or stay awake